10

15

## What is claimed is:

- 1. A process for preparing a LH-RH derivative which comprises subjecting a solution containing the LH-RH derivative to a step for treatment with a methacrylic synthetic adsorption resin and a step for treatment with an aromatic synthetic adsorption resin.
- 2. The process according to claim 1, wherein the LH-RH derivative is a peptide represented by the formula

5-oxo-Pro-His-Trp-Ser-Tyr-Y-Leu-Arg-Pro-Z

wherein Y indicates a residue selected from DLeu, DAla, DTrp, DSer(tBu), D2Nal and DHis(ImBzl), and Z indicates NH- $C_2H_5$  or Gly-NH<sub>2</sub>, respectively, or a salt thereof.

3. The process according to claim 1, wherein the LH-RH derivative is a peptide represented by the formula

5-oxo-Pro-His-Trp-Ser-Tyr-DLeu-Leu-Arg-Pro-NH- $C_2H_5$  or its acetate.

4. The process according to claim 1, wherein said process comprises using a methacrylic synthetic adsorption resin having a repeating unit represented by the formula

15

20

- 5. The process according to claim 1, wherein the aromatic synthetic adsorption resin is a styrene-divinylbenzene synthetic adsorption resin.
- 6. The process according to claim 5, wherein an average particle size of the styrene-divinylbenzene, synthetic adsorption resin is about 60  $\mu m$  to about 150  $\mu m$ .
- 7. The process according to claim 1, wherein said process comprises subjecting a solution containing the LH-RH derivative to the step for treatment with a methacrylic synthetic adsorption resin below about 10°C.
- 8. The process according to claim 1, wherein said process comprises subjecting a solution containing the LH-RH derivative to the step for treatment with an aromatic synthetic adsorption resin at about 10°C to about 20°C.
- 9. The process according to claim 1, wherein said process comprises subjecting a solution containing the LH-RH derivative to the step for treatment with a methacrylic, synthetic adsorption resin, followed by

10

15

20

25

subjecting to the step for treatment with an aromatic, synthetic adsorption resin.

- 10. The process according to claim 1, said process comprises passing a solution containing the LH-RH derivative through a resin in the step for treatment with a methacrylic synthetic adsorption resin and then eluting the LH-RH derivative, which is adsorbed on the resin, with an aqueous solution of acetic acid.
- 11. The process according to claim 10, wherein the concentration of an aqueous solution of acetic acid is about  $0.01~\mathrm{M}$  to about  $0.50~\mathrm{M}$ .
- 12. The process according to claim 1, wherein said process comprises passing a solution containing the LH-RH derivative through a resin in the step for treatment with a methacrylic, synthetic adsorption resin, followed by washing with an aqueous solution of ethanol, and then by eluting the LH-RH derivative that is adsorbed on the resin.
- 13. The process according to claim 1, wherein a solution containing the LH-RH derivative is that obtained by subjecting the LH-RH derivative protected with protective group(s) to a deprotection reaction followed by a neutralization reaction below about 10°C.
- 14. The process according to claim 1, wherein a solution containing the LH-RH derivative is that obtained by subjecting the LH-RH derivative protected with

5

protective group(s) to a deprotection reaction and then a neutralization reaction below about 10°C, followed by subjecting the resulting mixture to extraction of the LH-RH derivative and then concentration of the extract below 25°C.

The process according to claim 13 or 14, wherein the LH-RH derivative protected with protective group(s) is represented by t/te formula

5-oxo-Pro-His-Trp-Ser-Tyr-Y-Leu-Arg(X)-Pro-Z

wherein X indicates a protective group, Y indicates a residue selected from DLew, NAla, DTrp, DSer(tBu), D2Nal and DHis(ImBzl) Z indicates  $NH-C_2H_5$  or  $Gly-NH_2$ , and respectively.

- 16. Purified leuprorelin or a salt wherein the content of total related substances is about 1% or less.
- Purified leuprorelin or a salt thereof, wherein the content of 5-oxo-Pro-D-His-Trp-Ser-Tyr-D-Leu-Leu-Arg-Pro-NH-CH2-CH3 or a salt thereof is about 0.3% or less.

15